

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Editors: **W. Clegg** and **D. G. Watson**

Chloro{ μ -2-[(*E*)-1-(2-oxido-3-methylphenyl)ethylideneamino]-acetato}pentaphenylditin(IV)

Anthony Linden, Tushar S. Basu Baul and Cheerfulman Masharing

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

Anthony Linden,^{a*}
Tushar S. Basu Baul^b and
Cheerfulman Masharing^b^aInstitute of Organic Chemistry, University of
Zürich, Winterthurerstrasse 190, CH-8057
Zürich, Switzerland, and ^bDepartment of
Chemistry, North-Eastern Hill University,
NEHU Permanent Campus, Umshing,
Shillong 793 022, India

Correspondence e-mail: alinden@oci.unizh.ch


Key indicators

Single-crystal X-ray study
T = 160 K
Mean $\sigma(\text{C}-\text{C})$ = 0.009 Å
R factor = 0.045
wR factor = 0.113
Data-to-parameter ratio = 23.6For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.Chloro[μ -2-[(*E*)-1-(2-oxido-3-methylphenyl)-
ethylideneamino]acetato]pentaphenylditin(IV)

The title compound, $[\text{Sn}_2(\text{C}_6\text{H}_5)_5(\text{C}_{11}\text{H}_{11}\text{NO}_3)\text{Cl}]$, is a dinuclear organotin adduct in which the two Sn atoms are bridged *via* the carboxylate O—C—O group of a 2-[(*E*)-1-(2-hydroxyaryl)alkylideneamino]acetate ligand. Each Sn atom has a distorted trigonal bipyramidal geometry, with the Ph_3SnCl moiety being less distorted.


Comment

The title compound, (I), was prepared during an ongoing study of the coordination chemistry of organotin(IV) complexes of 2-[(*E*)-1-(2-hydroxyaryl)alkylideneamino]acetates (L). These ligand systems generate a great variety of structural forms with $R_2\text{Sn}$ - and $R_3\text{Sn}$ - moieties (Dakternieks *et al.*, 1998; Basu Baul & Tiekink, 1999; Basu Baul *et al.*, 2001, 2002, 2003, 2005). A few examples of dinuclear organotin adducts of the type $R_2\text{SnL}\cdot R_2\text{SnCl}_2$ ($R = \text{Ph}$, ^tBu; Khoo *et al.*, 1997; Dakternieks *et al.*, 1998) and $R_2\text{SnL}\cdot R_3\text{SnCl}$ ($R = \text{Ph}$; Dakternieks *et al.*, 1998; Basu Baul *et al.*, 2003) are known where two Sn atoms are bridged *via* the carboxylate O—C—O group of an L ligand. These considerations stirred our interest in the synthesis and structure of the title compound, (I), which has the $R_2\text{SnL}\cdot R_3\text{SnCl}$ ($R = \text{Ph}$) formulation.



The structure of (I) is virtually isomorphous with that of the $\text{Ph}_2\text{SnL}\cdot\text{Ph}_3\text{SnCl}$ adduct reported by Dakternieks *et al.* (1998). The only difference between the two compounds is the addition of the 3-methyl group on the benzene ring of the acetate ligand in (I). In all other respects, the two compounds and structures are the same and have similar coordination geometry at each Sn atom (Table 1). In (I), atom Sn1 has a distorted trigonal bipyramidal coordination geometry, with atoms O1 and O3 occupying axial positions and the O1—Sn—O3 angle distorted from linearity by 19.13 (13)°. The C10—Sn1—C11 angle is also about 14° wider than in an ideal trigonal bipyramid. Atom Sn1 lies 0.027 (1) Å out of the trigonal plane formed by atoms N1, C10 and C11 in the direction of atom O3. The geometry about atom Sn2 is also

Received 9 February 2005
Accepted 10 February 2005
Online 19 February 2005

**Figure 1**

View of the molecule of (I), showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 50% probability level and H atoms have been omitted for clarity.

distorted trigonal bipyramidal, with atoms Cl and O2 defining the axial positions, but the O1—Sn—O3 angle is distorted from linearity by only 3.76 (9)°. Atom Sn2 lies 0.224 (1) Å out of the trigonal plane formed by atoms C24, C30 and C36 in the direction of the Cl atom. The O2—Sn bond is about 0.24 Å longer than the O1—Sn1 bond and the carboxylate C—O distances are inversely related. The carboxylate C—O bonds are not completely delocalized, with the C1—O2 bond having much more double-bond character than the C1—O1 bond.

Experimental

Ph₂SnL was prepared by reacting Ph₂SnCl₂ and LHK as described by Basu Baul *et al.* (2001). A hot anhydrous benzene solution (10 ml) of Ph₃SnCl (0.20 g, 0.51 mmol) was added dropwise to a hot stirred benzene solution (20 ml) containing Ph₂SnL (0.25 g, 0.52 mmol) and refluxed for 1 h. The volatiles were removed *in vacuo*, the yellow mass was washed several times with hexane and filtered. The dried residue was dissolved in benzene and filtered to remove any particles. The filtrate was allowed to evaporate at room temperature, which afforded yellow crystals of (I) (52% yield, m.p. 457–458 K). Analysis calculated for C₄₁H₃₆ClNO₃Sn₂: C 57.03, H 4.20, N 1.62%; found: C 57.10, H 4.10, N 1.60%; IR (KBr, cm⁻¹): 1626 ν(OCO)_{asym}, 1235 ν[Ph(C=O)]; ¹H NMR (CDCl₃, 250.13 MHz): δ 7.83 [*m*, 4H, Sn—Ph^a(*ortho*-)], 7.67 [*m*, 6H, Sn—Ph^b(*ortho*-)], 7.48–7.35 [*m*, 17H, Sn—Ph^{a,b}(*meta*- and *para*-), H-7 and H-9], 6.70 (*t*, 1H, H-8), 4.29 (*s*, 2H, H-2), 2.63 (*s*, 3H, H-3'), 2.45 (*s*, 3H, H-6'); ¹³C NMR (CDCl₃, 62.89 MHz): δ 182.4 (C-1), 170.6 (C-3), 164.8 (C-5), 137.8 [Sn—Ph^a(*ipso*-)], 137.5 [Sn—Ph^b(*ipso*-)], 136.7 (C-7), 136.4 [Sn—Ph^a(*ortho*-)], 136.1 [Sn—Ph^b(*ortho*-)], 131.5 (C-9), 130.7 [Sn—Ph^a(*para*-)], 130.4 [Sn—Ph^b(*para*-)], 129.1 [Sn—Ph^b(*meta*-)], 128.9 [Sn—Ph^a(*meta*-)], 128.6 (C-4), 119.4 (C-6), 117.4 (C-8), 53.6 (C-2), 23.0 (C-3'), 16.8 (C-6') (*a* and *b* represent signals due to Sn—Ph₂ and Sn—Ph₃, respectively, and refer to the scheme for ligand assignment); ¹¹⁹Sn NMR (CDCl₃, 89.12 MHz): δ -51, -351. The ¹¹⁹Sn NMR data indicate the presence of two tin centres both having trigonal bipyramidal geometry in the structure (Dakternieks *et al.*, 1998; Basu Baul *et al.*, 2003). X-ray diffraction quality crystals were grown by slow evaporation of a solution of (I) in acetone.

Crystal data

[Sn₂(C₆H₅)₅(C₁₁H₁₁NO₃)Cl]
 $M_r = 863.39$
 Triclinic, $P\bar{1}$
 $a = 11.6984$ (2) Å
 $b = 13.0662$ (2) Å
 $c = 13.9410$ (3) Å
 $\alpha = 111.5066$ (9)°
 $\beta = 90.6470$ (9)°
 $\gamma = 115.4904$ (9)°
 $V = 1752.94$ (6) Å³

$Z = 2$
 $D_x = 1.636$ Mg m⁻³
 Mo K α radiation
 Cell parameters from 47226 reflections
 $\theta = 2.0$ – 30.0 °
 $\mu = 1.54$ mm⁻¹
 $T = 160$ (1) K
 Prism, yellow
 $0.15 \times 0.15 \times 0.08$ mm

Data collection

Nonius KappaCCD area-detector diffractometer
 φ and ω scans with κ offsets
 Absorption correction: multi-scan (Blessing, 1995)
 $T_{\min} = 0.759$, $T_{\max} = 0.881$
 57965 measured reflections

10272 independent reflections
 8257 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.060$
 $\theta_{\max} = 30.1$ °
 $h = -16 \rightarrow 16$
 $k = -18 \rightarrow 18$
 $l = -19 \rightarrow 19$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.045$
 $wR(F^2) = 0.113$
 $S = 1.20$
 10271 reflections
 436 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + 10.8098P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.29$ e Å⁻³
 $\Delta\rho_{\min} = -1.06$ e Å⁻³
 Extinction correction: SHELXL97
 Extinction coefficient: 0.0045 (3)

Table 1

Selected geometric parameters (Å, °).

Sn1—O3	2.057 (3)	Sn2—C30	2.132 (4)
Sn1—C11	2.116 (4)	Sn2—C24	2.153 (5)
Sn1—C10	2.119 (4)	Sn2—O2	2.415 (3)
Sn1—N1	2.160 (4)	Sn2—Cl	2.4610 (13)
Sn1—O1	2.172 (3)	O1—C1	1.276 (5)
Sn2—C36	2.117 (5)	O2—C1	1.239 (5)
O3—Sn1—C11	91.89 (15)	C30—Sn2—C24	111.18 (18)
O3—Sn1—C10	94.61 (15)	C36—Sn2—O2	83.24 (16)
C11—Sn1—C10	134.26 (17)	C30—Sn2—O2	84.30 (14)
O3—Sn1—N1	84.27 (13)	C24—Sn2—O2	84.51 (16)
C11—Sn1—N1	112.60 (15)	C36—Sn2—Cl	93.48 (14)
C10—Sn1—N1	113.09 (16)	C30—Sn2—Cl	96.33 (13)
O3—Sn1—O1	160.87 (13)	C24—Sn2—Cl	98.69 (15)
C11—Sn1—O1	94.66 (15)	O2—Sn2—Cl	176.24 (9)
C10—Sn1—O1	93.64 (15)	Cl—O1—Sn1	117.5 (3)
N1—Sn1—O1	76.61 (13)	Cl—O2—Sn2	134.2 (3)
C36—Sn2—C30	128.01 (18)	C5—O3—Sn1	128.3 (3)
C36—Sn2—C24	117.50 (18)		

The methyl groups were constrained to an ideal geometry (C—H = 0.98 Å), with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$, but were allowed to rotate freely about the parent C—C bonds. All other H atoms were placed in geometrically idealized positions and constrained to ride on their parent atoms, with C—H distances in the range 0.95–0.99 Å and with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$. One low angle reflection was partially obscured by the beam stop and was omitted from the refinement. The largest peak of residual electron density is 1.02 Å from atom H15, but is merely the highest peak in a high background of noise in the difference Fourier map. The deepest electron-density hole lies 0.80 Å from atom Sn2.

Data collection: COLLECT (Nonius, 2000); cell refinement: DENZO-SMN (Otwinowski & Minor, 1997); data reduction: DENZO-SMN and SCALEPACK (Otwinowski & Minor, 1997); program(s) used to solve structure: SHELXS97 (Sheldrick, 1997); program(s) used to refine structure: SHELXL97 (Sheldrick, 1997);

molecular graphics: *ORTEP* (Johnson, 1976); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2003).

The financial support (grant No. SP/S1/F26/99, TSBB) of the Department of Science & Technology, New Delhi, India, is gratefully acknowledged.

References

- Basu Baul, T. S., Dutta, S., Masharing, C., Rivarola, E. & Englert, U. (2003). *Heteroatom Chem.* **14**, 149–154.
- Basu Baul, T. S., Dutta, S., Rivarola, E., Butcher, R. & Smith, F. E. (2002). *J. Organomet. Chem.* **654**, 100–108.
- Basu Baul, T. S., Dutta, S., Rivarola, E., Scopelliti, M. & Choudhuri, S. (2001). *Appl. Organomet. Chem.* **15**, 947–953.
- Basu Baul, T. S., Masharing, C., Willem, R., Biesemans, M., Holčapek, M., Jirásko, R. & Linden, A. (2005). *J. Organomet. Chem.* Submitted.
- Basu Baul, T. S. & Tiekink, E. R. T. (1999). *Z. Kristallogr. New Cryst. Struct.* **214**, 361–362.
- Blessing, R. H. (1995). *Acta Cryst.* **A51**, 33–38.
- Dakternieks, D., Basu Baul, T. S., Dutta, S. & Tiekink, E. R. T. (1998). *Organometallics*, **17**, 3058–3062.
- Johnson, C. K. (1976). *ORTEP*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Khoo, L. E., Xu, Y., Goh, N. K., Chia, L. S. & Koh, L. L. (1997). *Polyhedron*, **16**, 573–576.
- Nonius (2000). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXL97* and *SHELXS97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.